arXiv:nlin/0004013v1 [nlin.SI] 9 Apr 2000

The critical $A^{(1)}_{n-1}$ chain

T. Kojima* and S. Yamasita†

*Department of Mathematics, College of Science and Technology, Nihon University,
Chiyoda-ku, Tokyo 101-0062, Japan
† Information Technology Center, Daiwa Institute of Research Ltd.,
Kōtō-ku, Tokyo 135-8460, Japan

February 8, 2008

Abstract

We study the $A^{(1)}_{n-1}$ spin chain at the critical regime $|q|=1$. We give the free boson realizations of the type-I vertex operators and their duals. Using these free boson realizations, we give the integral representations for the correlation functions.

1 Introduction

The one dimensional spin $A^{(1)}_{n-1}$ chain is described by the Hamiltonian :

$$\mathcal{H}=\sum_{k=-\infty}^{\infty}\left\{q\sum_{\substack{a,b=0\\a>b}}^{n-1}e_{aa}^{(k+1)}e_{bb}^{(k)}+q^{-1}\sum_{\substack{a,b=0\\a<b}}^{n-1}e_{aa}^{(k+1)}e_{bb}^{(k)}-\sum_{\substack{a,b=0\\a\neq b}}^{n-1}e_{ab}^{(k+1)}e_{ba}^{(k)}\right\}, \tag{1.1}$$

where $e_{ab}^{(k)}$ is a matrix unit acting on k th site. In this paper we consider the critical regime $|q|=1$. For the special case $n=2$ this model becomes the celebrated XXZ spin. M.Jimbo,H.Konno and T.Miwa [1] established the trace construction of the correlation functions for the XXZ chain at critical regime $|q|=1$. In this paper we give the correlation functions for the higher rank generalization of the XXZ chain at the critical regime $|q|=1$. In order to write down the correlation functions, we need the 'dual' vertex operators. This problem was absent from the XXZ chain, because the vertex operators are self-dual in this case. We give the pair of the vertex operators and their duals in this paper.

We give the N point correlation functions which describe the ground-state average $\langle\mathcal{O}\rangle$ of the local operator :

$$\mathcal{O}=e^{(1)}_{\epsilon_1\epsilon_1'}\cdots e^{(N)}_{\epsilon_N\epsilon_N'}. \tag{1.2}$$

The correlation functions of the critical $A_{n-1}^{(1)}$ chain are described by the following systems of difference equations :

$$\begin{aligned}
& G^{(N)}(\beta_1,\cdots,\beta_j-\lambda i,\cdots,\beta_N|\beta_{N+1},\cdots,\beta_{2N}) \\
= \ & R_{j,j-1}^{V^*V^*}(\beta_j-\beta_{j-1}-\lambda i)\cdots R_{j,1}^{V^*V^*}(\beta_j-\beta_1-\lambda i)R_{j,2N}^{V^*V}(\beta_j-\beta_{2N})\cdots R_{j,N+1}^{V^*V}(\beta_j-\beta_{N+1}) \\
\times \ & R_{j,N}^{V^*V^*}(\beta_j-\beta_N)\cdots R_{j,j+1}^{V^*V^*}(\beta_j-\beta_{j+1})G^{(N)}(\beta_1,\cdots,\beta_j,\cdots,\beta_N|\beta_{N+1},\cdots,\beta_{2N}),
\end{aligned} \tag{1.3}$$

and

$$\begin{aligned}
& G^{(N)}(\beta_1,\cdots,\beta_N|\beta_{N+1},\cdots,\beta_j+i\lambda,\cdots\beta_{2N}) \\
= \ & R_{j+1,j}^{VV}(\beta_{j+1}-\beta_j+\lambda i)\cdots R_{2N.j}^{VV}(\beta_{2N}-\beta_j+\lambda i)R_{1,j}^{V^*V}(\beta_1-\beta_j)\cdots R_{N,j}^{V^*V}(\beta_N-\beta_j) \\
\times \ & R_{N+1,j}^{VV}(\beta_{N+1}-\beta_j)\cdots R_{j-1,j}^{VV}(\beta_{j-1}-\beta_j)G^{(N)}(\beta_1,\cdots,\beta_N|\beta_{N+1},\cdots,\beta_j,\cdots\beta_{2N}).
\end{aligned} \tag{1.4}$$

Here $R_{ij}^{VV}(\beta) \in \mathrm{End}(V^{\otimes 2N})$ signifies the matrix acting as $R^{VV}(\beta)$ on th $(i,j)-$th tensor components and as identity elsewhere. Here $R^{VV}(\beta), R^{V^*V^*}(\beta)$ and $R^{V^*V}(\beta)$ are given by (3.1), (3.2) and (3.4). The correlation functions satisfy the restriction equation :

$$\begin{aligned}
& G^{(N)}(\beta+i(\pi-\lambda),\beta_2,\cdots,\beta_N|\beta_{N+1},\cdots,\beta_{2N})_{\epsilon_1\cdots\epsilon_N,\epsilon'_N\cdots\epsilon'_1} \\
= \ & \delta_{\epsilon_1,\epsilon'_1}G^{(N-1)}(\beta_2,\cdots,\beta_N|\beta_{N+1},\cdots,\beta_{2N-1})_{\epsilon_2\cdots\epsilon_N,\epsilon'_N\cdots\epsilon'_2}.
\end{aligned} \tag{1.5}$$

Here we have set

$$\begin{aligned}
& G^{(N)}(\beta_1\cdots\beta_N|\beta_{N+1}\cdots\beta_{2N}) \\
= \ & \sum_{\epsilon_1\cdots\epsilon_{2N}=0}^{n-1} v_{\epsilon_1}\otimes\cdots\otimes v_{\epsilon_{2N}}G^{(N)}(\beta_1\cdots\beta_N|\beta_{N+1}\cdots\beta_{2N})_{\epsilon_1\cdots\epsilon_{2N}}.
\end{aligned} \tag{1.6}$$

In this paper we set the deformation parameter as

$$q=-\exp\left(\frac{\pi i}{\xi}\right), \tag{1.7}$$

where $\xi > 1$.
The ground state averages are obtaind from the components $G^{(N)}$, by taking $\lambda = 2\pi$, and specializing the spectral parameters :

$$G^{(N)}(\beta+\pi i,\cdots,\beta+\pi i|\beta,\cdots,\beta)_{\epsilon_1\cdots\epsilon_N,\epsilon_N\cdots\epsilon_1}, \tag{1.8}$$

When we solve the diffence equations (1.3), (1.4), directly [2], we have a difficulty. The difficulty in this approach is that the solutions are determined only up to arbitrary periodic functions, so one has to single out in some way the correct solutions which correspond to the correlation functions. When we construct the solutions by the trace of the vertex operators, the ambiguity of solutions are resolved. In this paper we give the free boson realizations of the type-I vertex operators and their duals, and give the trace construction for the correlation functions of the critical $A_{n-1}^{(1)}$ spin. In this connection, we should mention about the work [3], in which the authors give the free boson realizations of the dual type-II vertex operators of the $A_{n-1}^{(1)}$ Toda field theory with imaginary coupling.

Now a few words about the organization of the paper. In section 2 we review the model briefly. In section 3 we give the defining relations of the vertex operators. In section 4 we give the free boson realizations of the vertex operators. In section 5 we give the proofs of the properties of the vertex operators. In section 6 we give the integral representations of the correlation functions. In Appendix A we summarized the multi Gamma functions. In Appendix B we summarized the normal ordering of the basic operators.

2 The critical $A_{n-1}^{(1)}$ chain

Let us set the R-matrix by

$$R^{VV}(\beta)=r(\beta)\bar{R}(\beta),\quad r(\beta)=-\frac{S_2(i\beta|\frac{2\pi}{n}\xi,2\pi)S_2(-i\beta+\frac{2\pi}{n}|\frac{2\pi}{n}\xi,2\pi)}{S_2(-i\beta|\frac{2\pi}{n}\xi,2\pi)S_2(i\beta+\frac{2\pi}{n}|\frac{2\pi}{n}\xi,2\pi)},\tag{2.1}$$

where $S_2(\beta|\omega_1,\omega_2)$ is the Multi-Sine function given in Appendix A.
The matrix $\bar{R}(\beta)$ is given as follows :

$$\bar{R}(\beta)v_{k_1}\otimes v_{k_2}=\sum_{j_1,j_2=0}^{n-1}v_{j_1}\otimes v_{j_2}\bar{R}(\beta)_{j_1j_2}^{k_1k_2},\tag{2.2}$$

where the nonzero entrise are

$$\begin{aligned}\bar{R}(\beta)_{jj}^{jj} &= 1, &(2.3)\\ \bar{R}(\beta)_{jk}^{jk} &= -\frac{\mathrm{sh}\left(\frac{n}{2\xi}\beta\right)}{\mathrm{sh}\left(\frac{n}{2\xi}(\beta+\frac{2\pi i}{n})\right)},\quad (j\neq k), &(2.4)\\ \bar{R}(\beta)_{jk}^{kj} &= \begin{cases}\dfrac{e^{-\frac{n}{2\xi}\beta}\mathrm{sh}\left(\frac{\pi i}{\xi}\right)}{\mathrm{sh}\left(\frac{n}{2\xi}(\beta+\frac{2\pi i}{n})\right)}, & (j<k),\\ \dfrac{e^{\frac{n}{2\xi}\beta}\mathrm{sh}\left(\frac{\pi i}{\xi}\right)}{\mathrm{sh}\left(\frac{n}{2\xi}(\beta+\frac{2\pi i}{n})\right)}, & (j>k),\end{cases} &(2.5)\end{aligned}$$

The R-matrix satisfies the Yang-Baxter equations,

$$R_{12}^{VV}(\beta_1-\beta_2)R_{13}^{VV}(\beta_1-\beta_3)R_{23}^{VV}(\beta_2-\beta_3)=R_{23}^{VV}(\beta_2-\beta_3)R_{13}^{VV}(\beta_1-\beta_3)R_{12}^{VV}(\beta_1-\beta_2),\tag{2.6}$$

and the unitarity,

$$R_{12}^{VV}(\beta1-\beta_2)R_{21}^{VV}(\beta_2-\beta_1)=id.\tag{2.7}$$

Let us set the monodromy matrix $\mathcal{T}(\beta)$ acting on the $(N+1)-$ fold tensor product $V_0\otimes V_1\otimes\cdots\otimes V_N$, $(V_k = V = \mathbb{C}^n)$,

$$\mathcal{T}(\beta) = R_{0,1}^{VV}(\beta)\cdots R_{0,N}^{VV}(\beta) = \begin{pmatrix} A_{1,1} & A_{1,2} & \cdots & A_{1,n-1} & A_{1,n} \\ A_{2,1} & A_{2,2} & \cdots & A_{2,n-1} & A_{2,n} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ A_{n-1,1} & A_{n-1,2} & \cdots & A_{n-1,n-1} & A_{n-1,n} \\ A_{n,1} & A_{n,2} & \cdots & A_{n,n-1} & A_{n,n} \end{pmatrix}, \tag{2.8}$$

where the partition into $n\times n$ blocks is according to the base of V_0.
Let us set the transfer matrix $T(\beta)$ acting on the N-th fold tensor product $V_1\otimes\cdots\otimes V_N$ by

$$T(\beta) = \sum_{j=1}^{n} A_{j,j} = \mathrm{tr}_{V_0}(\mathcal{T}(\beta)). \tag{2.9}$$

From the Yang-Baxter equation, we know the transfer matrices commute each other.

$$[T(\beta_1), T(\beta_2)] = 0. \tag{2.10}$$

In the thermodynamic limit $N\to\infty$, the logarithmic derivative of the transfer matrix and the Hamiltonian (1.1) have the following relation.

$$\left(\frac{d}{d\beta}\mathrm{log}T\right)(0) \sim \mathcal{H}. \tag{2.11}$$

A.Doikou and R.I.Nepomechie [5] computed by Bethe Ansatz the scattering matrix for the critical $A_{n-1}^{(1)}$ spin chain. The scattering matrix $S(\beta)$ is given by

$$S(\beta) = s(\beta)\bar{S}(\beta), \quad s(\beta) = \frac{S_2(-i\beta|\frac{2\pi}{n}(\xi-1), 2\pi)S_2(i\beta+\frac{2(n-1)\pi}{n}|\frac{2\pi}{n}(\xi-1), 2\pi)}{S_2(i\beta|\frac{2\pi}{n}(\xi-1), 2\pi)S_2(-i\beta+\frac{2(n-1)\pi}{n}|\frac{2\pi}{n}(\xi-1), 2\pi)}. \tag{2.12}$$

The matrix $\bar{S}(\beta)$ is given as follows :

$$\bar{S}(\beta)v_{k_1}\otimes v_{k_2} = \sum_{j_1,j_2=0}^{n-1} v_{j_1}\otimes v_{j_2}\bar{S}(\beta)_{j_1j_2}^{k_1k_2}, \tag{2.13}$$

where the nonzero entrise are

$$\bar{S}(\beta)_{jj}^{jj} = 1, \tag{2.14}$$

$$\bar{S}(\beta)_{jk}^{jk} = -\frac{\mathrm{sh}\left(\frac{n}{2(\xi-1)}\beta\right)}{\mathrm{sh}\left(\frac{n}{2(\xi-1)}(\beta-\frac{2\pi i}{n})\right)}, \quad (j\neq k), \tag{2.15}$$

$$\bar{S}(\beta)_{jk}^{kj} = \begin{cases} -\dfrac{e^{-\frac{n}{2(\xi-1)}\beta}\mathrm{sh}\left(\frac{\pi i}{(\xi-1)}\right)}{\mathrm{sh}\left(\frac{n}{2(\xi-1)}(\beta-\frac{2\pi i}{n})\right)}, & (j<k), \\ -\dfrac{e^{\frac{n}{2(\xi-1)}\beta}\mathrm{sh}\left(\frac{\pi i}{(\xi-1)}\right)}{\mathrm{sh}\left(\frac{n}{2(\xi-1)}(\beta-\frac{2\pi i}{n})\right)}, & (j>k), \end{cases} \tag{2.16}$$

The scattering matrix of the present model agrees with the scattering matrix of the $A_{n-1}^{(1)}$ Toda fields theory with imaginary coupling.

3 Vertex Operators

The type-I vertex operators satisfy

$$\Phi_{j_2}(\beta_1)\Phi_{j_1}(\beta_2) = \sum_{k_1 k_2=0}^{n-1} R^{VV}(\beta_1-\beta_2)_{j_1 j_2}^{k_1 k_2}\Phi_{k_1}(\beta_2)\Phi_{k_2}(\beta_1). \tag{3.1}$$

The dual type-I vertex operators satisfy

$$\Phi_{j_2}^*(\beta_1)\Phi_{j_1}^*(\beta_2) = \sum_{k_1 k_2=0}^{n-1} R^{V^*V^*}(\beta_1-\beta_2)_{j_1 j_2}^{k_1 k_2}\Phi_{k_1}^*(\beta_2)\Phi_{k_2}^*(\beta_1), \tag{3.2}$$

where we set

$$R^{V^*V^*}(\beta)_{j_1 j_2}^{k_1 k_2} = R^{VV}(\beta)_{j_2 j_1}^{k_2 k_1}. \tag{3.3}$$

The type-I and dual type-I vertex operators satisfy

$$\Phi_{j_2}(\beta_1)\Phi_{j_1}^*(\beta_2) = \sum_{k_1 k_2=0}^{n-1} R^{V^*V}(\beta_1-\beta_2)_{j_1 j_2}^{k_1 k_2}\Phi_{k_1}^*(\beta_2)\Phi_{k_2}(\beta_1), \tag{3.4}$$

where we set

$$R^{VV^*}(\beta) = r^*(\beta)\bar{R}^*(\beta), \quad r^*(\beta) = \frac{S_2(-i\alpha+\pi|\frac{2\pi}{n}\xi,2\pi)S_2(i\alpha+\pi+\frac{2\pi}{n}|\frac{2\pi}{n}\xi,2\pi)}{S_2(i\alpha+\pi|\frac{2\pi}{n}\xi,2\pi)S_2(-i\alpha+\pi+\frac{2\pi}{n}|\frac{2\pi}{n}\xi,2\pi)}. \tag{3.5}$$

The matrix $\bar{R}^*(\beta)$ is given as follows :

$$\bar{R}^*(\beta)v_{k_1}\otimes v_{k_2} = \sum_{j_1,j_2=0}^{n-1} v_{j_1}\otimes v_{j_2}\bar{R}^*(\beta)_{j_1 j_2}^{k_1 k_2}, \tag{3.6}$$

where the nonzero entrise are

$$\bar{R}^*(\beta)_{jk}^{jk} = 1, \quad (j\neq k), \tag{3.7}$$

$$\bar{R}^*(\beta)_{jj}^{jj} = -\frac{\mathrm{sh}\left(\frac{n}{2\xi}\beta\right)}{\mathrm{sh}\left(\frac{n}{2\xi}(\beta+\frac{2\pi i}{n})\right)}, \tag{3.8}$$

$$\bar{R}^*(\beta)_{jj}^{kk} = \begin{cases} \dfrac{e^{-\frac{n}{2\xi}\beta}\mathrm{sh}\left(\frac{\pi}{\xi}i\right)}{\mathrm{sh}\left(\frac{n}{2\xi}(\beta+\frac{2\pi i}{n})\right)}, & (j>k), \\ \dfrac{e^{\frac{n}{2\xi}\beta}\mathrm{sh}\left(\frac{\pi}{\xi}i\right)}{\mathrm{sh}\left(\frac{n}{2\xi}(\beta+\frac{2\pi i}{n})\right)}, & (j<k), \end{cases} \tag{3.9}$$

The type-I and the dual type-I vertex operators satisfy

$$\Phi_{j_2}^*(\beta_1)\Phi_{j_1}(\beta_2)=\sum_{k_1k_2=0}^{n-1}R^{VV^*}(\beta_1-\beta_2)_{j_1j_2}^{k_1k_2}\Phi_{k_1}(\beta_2)\Phi_{k_2}^*(\beta_1), \tag{3.10}$$

where we set

$$R^{VV^*}(\beta)_{j_1j_2}^{k_1k_2}=R^{V^*V}(\beta)_{k_1k_2}^{j_1j_2}. \tag{3.11}$$

The type-I vertex operator and it's dual satisfy the inversion relation :

$$\Phi_{j_1}(\beta)\Phi_{j_2}^*(\beta+\pi i)=g_n^{-1}e^{\frac{2\pi i}{n}j_1}\delta_{j_1,j_2}id, \quad (j_1\leq j_2). \tag{3.12}$$

where we set

$$g_n^{-1}=\left(\frac{2\xi}{n}\right)^{n-1}\mathrm{sh}\left(\frac{\pi i}{\xi}\right)e^{-\frac{\xi-1}{\xi}(\gamma+\log\frac{2\pi\xi}{n})n+\frac{\pi i}{n}(1-n)}\frac{\Gamma(1/\xi)^{n-1}}{\Gamma(1-1/\xi)}. \tag{3.13}$$

4 Free boson realizations

In this section we give the free boson realizations of the vertex operators.

Let us set bose-fields as

$$[b_j(t),b_k(t')]=-\frac{1}{t}\frac{\mathrm{sh}\left(\frac{\pi}{n}(a_j|a_k)t\right)\mathrm{sh}\left(\frac{\pi}{n}(\xi-1)t\right)}{\mathrm{sh}\left(\frac{\pi}{n}t\right)\mathrm{sh}\left(\frac{\pi}{n}\xi t\right)}\delta(t+t') \tag{4.1}$$

Let us set

$$b_1^*(t) = -\sum_{j=1}^{n-1}b_j(t)\frac{\mathrm{sh}\frac{(n-j)\pi t}{n}}{\mathrm{sh}\pi t}, \tag{4.2}$$

$$b_{n-1}^*(t) = -\sum_{j=1}^{n-1}b_j(t)\frac{\mathrm{sh}\frac{j\pi t}{n}}{\mathrm{sh}\pi t}. \tag{4.3}$$

We have

$$[b_1^*(t),b_j(t')] = \delta_{j1}\frac{1}{t}\frac{\mathrm{sh}\left(\frac{\pi}{n}(\xi-1)t\right)}{\mathrm{sh}\left(\frac{\pi}{n}\xi t\right)}\delta(t+t'), \tag{4.4}$$

$$[b_j(t),b_{n-1}^*(t')] = \delta_{j,n-1}\frac{1}{t}\frac{\mathrm{sh}\left(\frac{\pi}{n}(\xi-1)t\right)}{\mathrm{sh}\left(\frac{\pi}{n}\xi t\right)}\delta(t+t'). \tag{4.5}$$

Let us set the basic operators as

$$U_j(\beta) = :\exp\left(-\int_{-\infty}^{\infty}b_j(t)e^{i\beta t}dt\right): \ (1\leq j\leq n-1), \tag{4.6}$$

$$U_0(\beta) = :\exp\left(-\int_{-\infty}^{\infty}b_1^*(t)e^{i\beta t}dt\right):, \tag{4.7}$$

$$U_n(\beta) = :\exp\left(-\int_{-\infty}^{\infty}b_{n-1}^*(t)e^{i\beta t}dt\right):. \tag{4.8}$$

The bosonization of the type-I vertex operator is given by

$$\begin{aligned}\Phi_j(\beta) &= \int_{-\infty}^{\infty}\frac{d\alpha_1}{2\pi i}\cdots\int_{-\infty}^{\infty}\frac{d\alpha_j}{2\pi i}\frac{e^{\frac{n}{2\xi}(\alpha_j-\beta)}U_0(\beta)U_1(\alpha_1)\cdots U_j(\alpha_j)}{\prod_{k=1}^{j}\mathrm{sh}\left(\frac{n}{2\xi}(\alpha_{k-1}-\alpha_k-\frac{\pi i}{n})\right)} \qquad (4.9)\\ &= \int_{-\infty}^{\infty}\frac{d\alpha_1}{2\pi i}\cdots\int_{-\infty}^{\infty}\frac{d\alpha_j}{2\pi i}\frac{e^{\frac{n}{2\xi}(\alpha_j-\beta)}U_j(\alpha_j)U_{j-1}(\alpha_{j-1})\cdots U_0(\beta)}{\prod_{k=1}^{j}\mathrm{sh}\left(\frac{n}{2\xi}(\alpha_k-\alpha_{k-1}-\frac{\pi i}{n})\right)}, \ (0\leq j\leq n-1)\end{aligned}$$

where $\alpha_0 = \beta$.

The bosonization of the dual type-I vertex operator is given by

$$\begin{aligned}\Phi_j^*(\beta) &= \int_{-\infty}^{\infty} \frac{d\alpha_{j+1}}{2\pi i} \cdots \int_{-\infty}^{\infty} \frac{d\alpha_{n-1}}{2\pi i} \frac{e^{\frac{n}{2\xi}(\alpha_{j+1}-\beta)} U_{j+1}(\alpha_{j+1}) \cdots U_{n-1}(\alpha_{n-1}) U_n(\beta)}{\prod_{k=j+1}^{n-1} \mathrm{sh}\left(\frac{n}{2\xi}(\alpha_k - \alpha_{k+1} - \frac{\pi i}{n})\right)}, \qquad (4.10)\\ &= \int_{-\infty}^{\infty} \frac{d\alpha_{j+1}}{2\pi i} \cdots \int_{-\infty}^{\infty} \frac{d\alpha_{n-1}}{2\pi i} \frac{e^{\frac{n}{2\xi}(\alpha_{j+1}-\beta)} U_n(\beta) U_{n-1}(\alpha_{n-1}) \cdots U_{j+1}(\alpha_{j+1})}{\prod_{k=j+1}^{n-1} \mathrm{sh}\left(\frac{n}{2\xi}(\alpha_{k+1} - \alpha_k - \frac{\pi i}{n})\right)}, \ (0 \le j \le n-1),\end{aligned}$$

where $\alpha_n = \beta$.

T.Miwa and Y.Takeyama [3] gave the bosonization of some vertex operator, to construct the solutions of the critical quantum Knizhnik-Zamolodchikov equations at Level 0. The calculation of S-matrix of the critical $A_{n-1}^{(1)}$ chain [5] teaches us that their vertex operators are the dual type-II vertex operators of the present model.

5 Proof

In this section we prove that the bosonizations of the vertex operators satisfy the commutation relations (3.1), (3.2), (3.4), (3.10) and the inversion relation (3.12). For convenience we list below formulae for the contractions of the basic operators.

$$U_j(\beta_1) U_k(\beta_2) = U_k(\beta_2) U_j(\beta_1), \quad (|j-k| \ge 2). \qquad (5.1)$$

$$U_j(\beta_1) U_j(\beta_2) = U_j(\beta_2) U_j(\beta_1) r(\beta_1 - \beta_2), \quad (j = 0, n), \qquad (5.2)$$

$$r(\beta) = -\frac{S_2(i\alpha|\frac{2\pi}{n}\xi, 2\pi) S_2(-i\alpha + \frac{2\pi}{n}|\frac{2\pi}{n}\xi, 2\pi)}{S_2(-i\alpha|\frac{2\pi}{n}\xi, 2\pi) S_2(i\alpha + \frac{2\pi}{n}|\frac{2\pi}{n}\xi, 2\pi)}, \qquad (5.3)$$

$$U_0(\beta_1) U_{n-1}(\beta_2) = r^*(\beta_1 - \beta_2) U_{n-1}(\beta_2) U_0(\beta_1), \qquad (5.4)$$

$$r^*(\beta) = \frac{S_2(-i\alpha + \pi|\frac{2\pi}{n}\xi, 2\pi) S_2(i\alpha + \pi + \frac{2\pi}{n}|\frac{2\pi}{n}\xi, 2\pi)}{S_2(i\alpha + \pi|\frac{2\pi}{n}\xi, 2\pi) S_2(-i\alpha + \pi + \frac{2\pi}{n}|\frac{2\pi}{n}\xi, 2\pi)}. \qquad (5.5)$$

$$\begin{aligned} U_j(\beta_1) U_j(\beta_2) &= H(\beta_1 - \beta_2) U_j(\beta_2) U_j(\beta_1), \quad (1 \le j \le n-1), & (5.6)\\ H(\beta) &= -\frac{\mathrm{sh}\left(\frac{n}{2\xi}(\beta + \frac{2\pi i}{n})\right)}{\mathrm{sh}\left(\frac{n}{2\xi}(-\beta + \frac{2\pi i}{n})\right)}, & (5.7)\end{aligned}$$

$$\begin{aligned} U_j(\beta_1) U_{j-1}(\beta_2) &= I(\beta_1 - \beta_2) U_{j-1}(\beta_2) U_j(\beta_1), \quad (1 \le j \le n), & (5.8)\\ I(\beta) &= \frac{\mathrm{sh}\left(\frac{n}{2\xi}\left(\beta - \frac{\pi i}{n}\right)\right)}{\mathrm{sh}\left(\frac{n}{2\xi}\left(-\beta - \frac{\pi i}{n}\right)\right)}. & (5.9)\end{aligned}$$

We are to prove the various properties of the vertex operators along the line of the papers [7, 8]. We are to prove the commutation relation of the type-I vertex operators (3.1). Consider the integral of the form :

$$\int_{-\infty}^{\infty}\int_{-\infty}^{\infty} d\alpha_1 d\alpha_2 U_j(\alpha_1)U_j(\alpha_2)F(\alpha_1,\alpha_2). \tag{5.10}$$

Due to the commutation relation of $U_j(\alpha)$, the above integral equals to

$$\int_{-\infty}^{\infty}\int_{-\infty}^{\infty} d\alpha_1 d\alpha_2 U_j(\alpha_1)U_j(\alpha_2)F(\alpha_2,\alpha_1)H(\alpha_2-\alpha_1) \tag{5.11}$$

Observing this we define 'weak equality' in the following sense. We say that the fuctions $G_1(\alpha_1,\alpha_2)$ and $G_2(\alpha_1,\alpha_2)$ are equal in weak sense if

$$G_1(\alpha_1,\alpha_2)+H(\alpha_2-\alpha_1)G_1(\alpha_2,\alpha_1)=G_2(\alpha_1,\alpha_2)+H(\alpha_2-\alpha_1)G_2(\alpha_2,\alpha_1). \tag{5.12}$$

We write

$$G_1(\alpha_1,\alpha_2)\sim G_2(\alpha_1,\alpha_2). \tag{5.13}$$

To prove the commutation relations (3.1) and (3.2) it is enough to prove the equalities of the integrand parts in weakly sense.

Firsi we will show

$$\Phi_\mu(\alpha_0)\Phi_\mu(\alpha_0')=r(\alpha_0-\alpha_0')\Phi_\mu(\alpha_0')\Phi_\mu(\alpha_0) \quad (0\le\mu\le n-1). \tag{5.14}$$

In what follows we use the notations :

$$b(\alpha)=-\frac{\mathrm{sh}\rho\alpha}{\mathrm{sh}\rho(\alpha+\frac{2\pi i}{n})},\quad c(\alpha)=\frac{\mathrm{sh}\rho\frac{2\pi i}{n}}{\mathrm{sh}\rho(\alpha+\frac{2\pi i}{n})},\quad d(\alpha)=\frac{e^{\rho\alpha}}{\mathrm{sh}\rho(-\alpha-\frac{\pi i}{n})},\quad \rho=\frac{n}{2\xi}. \tag{5.15}$$

For $\mu=0$ case it is just the commutation relation of $U_0(\alpha)$. For $\mu\ge 1$ case we will show that by induction. By using the commutation relations of the basic operators, we can rearrange the operator part as

$$U_0(\alpha_0)U_0(\alpha_0')U_1(\alpha_1)U_1(\alpha_1')\cdots U_\mu(\alpha_\mu)U_\mu(\alpha_\mu'). \tag{5.16}$$

The integrand function of (LHS) of (5.14) is given by

$$L_\mu(\alpha_0\alpha_0'\cdots\alpha_\mu\alpha_\mu')=\prod_{k=1}^{\mu}d(\alpha_k-\alpha_{k-1})\prod_{k=1}^{\mu}d(\alpha_k'-\alpha_{k-1}')\prod_{k=1}^{\mu}I(\alpha_k-\alpha_{k-1}') \tag{5.17}$$

The integrand of (RHS) is given by the exchange of variables $\alpha_0\leftrightarrow\alpha_0'$ of (LHS).

$$R_\mu(\alpha_0\alpha_0'\cdots\alpha_\mu\alpha_\mu')=L_\mu(\alpha_0'\alpha_0\cdots\alpha_\mu\alpha_\mu'). \tag{5.18}$$

For $\mu=1$ case the weakely identity for variables α_1,α_1' can be shown by exact calculation.

$$L_1(\alpha_0\alpha_0'\alpha_1\alpha_1')\sim R_1(\alpha_0\alpha_0'\alpha_1\alpha_1'). \tag{5.19}$$

Because the integrand function splits into two parts :

$$L_\mu(\alpha_0\alpha'_0\alpha_1\alpha'_1\cdots\alpha_\mu\alpha'_\mu)=L_1(\alpha_0\alpha'_0\alpha_1\alpha'_1)L_{\mu-1}(\alpha_1\alpha'_1\alpha_2\alpha'_2\cdots\alpha_\mu\alpha'_\mu), \tag{5.20}$$

the case $\mu\geq 2$ follows from induction.

Next we will show

$$\begin{aligned}&\Phi_\mu(\alpha_0)\Phi_\nu(\alpha'_0)\\ =\;& r(\alpha_0-\alpha'_0)\left(b(\alpha_0-\alpha'_0)\Phi_\nu(\alpha'_0)\Phi_\mu(\alpha_0)+e^{\mathrm{sgn}(\nu-\mu)\rho(\alpha_0-\alpha'_0)}c(\alpha_0-\alpha'_0)\Phi_\mu(\alpha'_0)\Phi_\nu(\alpha_0)\right).\end{aligned} \tag{5.21}$$

We prove the case $\nu>\mu$. The case $\mu>\nu$ is similar. For the case $\nu>\mu=0$, the equality (5.21) follows from the following integrand equality, which can be derived by direct calculations.

$$d(\alpha'_1-\alpha'_0)=b(\alpha_0-\alpha'_0)I(\alpha'_1-\alpha_0)d(\alpha'_1-\alpha'_0)+c(\alpha_0-\alpha'_0)e^{\rho(\alpha_0-\alpha'_0)}d(\alpha'_1-\alpha_0). \tag{5.22}$$

For the case $\nu>\mu\geq 1$, the equality (5.21) follows from the weak equality with respect to the variables $(\alpha_1\alpha'_1),(\alpha_2\alpha'_2)\cdots(\alpha_\mu\alpha'_\mu)$:

$$A_\mu(\alpha_0\alpha'_0\cdots\alpha_{\mu+1}\alpha'_{\mu+1})+B_\mu(\alpha_0\alpha'_0\cdots\alpha_{\mu+1}\alpha'_{\mu+1})+C_\mu(\alpha_0\alpha'_0\cdots\alpha_{\mu+1}\alpha'_{\mu+1})\sim 0, \tag{5.23}$$

where we set

$$A_\mu(\alpha_0\alpha'_0\cdots\alpha_{\mu+1}\alpha'_{\mu+1})=\prod_{k=0}^{\mu-1}I(\alpha_{k+1}-\alpha'_k)\prod_{k=0}^{\mu-1}d(\alpha_{k+1}-\alpha_k)\prod_{k=0}^{\mu}d(\alpha'_{k+1}-\alpha'_k), \tag{5.24}$$

$$\begin{aligned}B_\mu(\alpha_0\alpha'_0\cdots\alpha_{\mu+1}\alpha'_{\mu+1})\;=\;&-b(\alpha_0-\alpha'_0)\times I(\alpha'_{\mu+1}-\alpha'_\mu)\left\{\prod_{k=1}^{\mu-1}d(\alpha_{k+1}-\alpha'_k)\right\}I(\alpha_1-\alpha_0)\\ \times\;& d(\alpha'_{\mu+1}-\alpha_\mu)\left\{\prod_{k=1}^{\mu-1}d(\alpha_{k+1}-\alpha_k)\right\}d(\alpha_1-\alpha'_0)\left\{\prod_{k=1}^{\mu-1}d(\alpha'_{k+1}-\alpha'_k)\right\}d(\alpha'_1-\alpha_0),\end{aligned} \tag{5.25}$$

and

$$\begin{aligned}C_\mu(\alpha_0\alpha'_0\cdots\alpha_{\mu+1}\alpha'_{\mu+1})\;=\;&-c(\alpha_0-\alpha'_0)e^{\rho(\alpha_0-\alpha'_0)}\times\left\{\prod_{k=1}^{\mu-1}I(\alpha_{k+1}-\alpha'_k)\right\}I(\alpha_1-\alpha_0)\\ \times\;&\left\{\prod_{k=1}^{\mu}d(\alpha'_{k+1}-\alpha'_k)\right\}d(\alpha'_1-\alpha_0)\left\{\prod_{k=1}^{\mu}d(\alpha_{k+1}-\alpha_k)\right\}d(\alpha_1-\alpha'_0).\end{aligned} \tag{5.26}$$

We are to prove the equation (5.23) by induction of μ. Inserting the equation (5.23) for $\mu-1$ to B_μ, we have we have the equation in weakly sense with respect to the variables $(\alpha_1\alpha'_1)\cdots(\alpha_\mu\alpha'_\mu)$:

$$B_\mu(\alpha_0\alpha'_0\cdots\alpha_{\mu+1}\alpha'_{\mu+1})\sim A'_\mu(\alpha_0\alpha'_0\cdots\alpha_{\mu+1}\alpha'_{\mu+1})+C'_\mu(\alpha_0\alpha'_0\cdots\alpha_{\mu+1}\alpha'_{\mu+1}), \tag{5.27}$$

where we set

$$\begin{aligned}A'_\mu(\alpha_0\alpha'_0\cdots\alpha_\mu\alpha'_{\mu+1})=-\frac{b(\alpha_0-\alpha'_0)}{b(\alpha'_1-\alpha_1)}\left\{\prod_{k=2}^{\mu}d(\alpha'_{k+1}-\alpha'_k)\right\}d(\alpha'_2-\alpha_1)d(\alpha_1-\alpha'_0)\\ \times\left\{\prod_{k=2}^{\mu-1}d(\alpha_{k+1}-\alpha_k)\right\}d(\alpha_2-\alpha'_1)d(\alpha'_1-\alpha_0)\left\{\prod_{k=2}^{\mu-1}I(\alpha_{k+1}-\alpha'_k)\right\}I(\alpha_2-\alpha_1)I(\alpha_1-\alpha_0),\end{aligned} \tag{5.28}$$

and

$$\begin{aligned} C'_\mu(\alpha_0\alpha'_0\cdots\alpha_{\mu+1}\alpha'_{\mu+1}) &= \frac{b(\alpha_0-\alpha'_0)c(\alpha'_1-\alpha_1)e^{\rho(\alpha'_1-\alpha_1)}}{b(\alpha'_1-\alpha_1)} \\ \times\ \left\{\prod_{k=1}^{\mu} d(\alpha'_{k+1}-\alpha'_k)\right\} d(\alpha'_1-\alpha_0) &\left\{\prod_{k=1}^{\mu-1} d(\alpha_{k+1}-\alpha_k)\right\} d(\alpha_1-\alpha'_0) \left\{\prod_{k=1}^{\mu-1} I(\alpha_{k+1}-\alpha'_k)\right\} I(\alpha_1-\alpha_0). \end{aligned} \tag{5.29}$$

Using the following relation : $b(\alpha)H(\alpha)=b(-\alpha)$, we have

$$A'_\mu(\alpha_0\alpha'_0\cdots\alpha_{\mu+1}\alpha'_{\mu+1}) \sim A''_\mu(\alpha_0\alpha'_0\cdots\alpha_{\mu+1}\alpha'_{\mu+1}), \tag{5.30}$$

where

$$\begin{aligned} & A''_\mu(\alpha_0\alpha'_0\cdots\alpha_{\mu+1}\alpha'_{\mu+1}) \\ = \ & -\frac{b(\alpha_0-\alpha'_0)}{b(\alpha'_1-\alpha_1)}\left\{\prod_{k=0}^{\mu} d(\alpha'_{k+1}-\alpha'_k)\right\}\left\{\prod_{k=0}^{\mu} d(\alpha_{k+1}-\alpha_k)\right\}\left\{\prod_{k=0}^{\mu-1} I(\alpha_{k+1}-\alpha'_k)\right\} I(\alpha'_1-\alpha_0). \end{aligned} \tag{5.31}$$

We have

$$\begin{aligned} A_\mu+A''_\mu \ \sim\ & \frac{\mathrm{sh}\left(\rho(\alpha'_0-\alpha'_1-\frac{\pi i}{n})\right)\mathrm{sh}\left(\rho(\alpha_0-\alpha_1+\frac{\pi i}{n})\right)\mathrm{sh}\left(\rho(\alpha_0+\alpha_1-\alpha'_0-\alpha'_1)\right)\mathrm{sh}\left(\rho(\frac{2\pi i}{n})\right)}{\mathrm{sh}\left(\rho(\alpha'_0-\alpha_1-\frac{\pi i}{n})\right)\mathrm{sh}\left(\rho(\alpha'_1-\alpha_0+\frac{2\pi i}{n})\right)\mathrm{sh}\left(\rho(\alpha_0-\alpha'_1-\frac{\pi i}{n})\right)\mathrm{sh}\left(\rho(\alpha_0-\alpha'_0+\frac{2\pi i}{n})\right)} \\ \times\ & \frac{1}{b(\alpha'_1-\alpha_1)}\left\{\prod_{k=0}^{\mu} d(\alpha'_{k+1}-\alpha'_k)\right\}\left\{\prod_{k=0}^{\mu-1} d(\alpha_{k+1}-\alpha_k)\right\}\left\{\prod_{k=1}^{\mu-1} I(\alpha_{k+1}-\alpha'_k)\right\}, \end{aligned} \tag{5.32}$$

where we have used the relation :

$$\begin{aligned} & I(\alpha_1-\alpha'_0)b(\alpha'_1-\alpha_1)-I(\alpha'_1-\alpha_0)b(\alpha_0-\alpha'_0) \\ = \ & \frac{\mathrm{sh}\left(\rho(\alpha'_0-\alpha'_1-\frac{\pi i}{n})\right)\mathrm{sh}\left(\rho(\alpha_0-\alpha_1+\frac{\pi i}{n})\right)\mathrm{sh}\left(\rho(\alpha_0+\alpha_1-\alpha'_0-\alpha'_1)\right)\mathrm{sh}\left(\rho(\frac{2\pi i}{n})\right)}{\mathrm{sh}\left(\rho(\alpha'_0-\alpha_1-\frac{\pi i}{n})\right)\mathrm{sh}\left(\rho(\alpha'_1-\alpha_0+\frac{2\pi i}{n})\right)\mathrm{sh}\left(\rho(\alpha_0-\alpha'_1-\frac{\pi i}{n})\right)\mathrm{sh}\left(\rho(\alpha_0-\alpha'_0+\frac{2\pi i}{n})\right)}. \end{aligned} \tag{5.33}$$

We have

$$\begin{aligned} C_\mu+C'_\mu &= -b(\alpha_0-\alpha'_0)\frac{\mathrm{sh}\left(\rho(\frac{2\pi i}{n})\right)\mathrm{sh}\left(\rho(\alpha_0+\alpha_1-\alpha'_0-\alpha'_1))\right)}{\mathrm{sh}\left(\rho(\alpha'_1-\alpha_1)\right)\mathrm{sh}\left(\rho(\alpha_0-\alpha'_0))\right)} \\ \left\{\prod_{k=1}^{\mu} d(\alpha'_{k+1}-\alpha'_k)\right\} d(\alpha'_1-\alpha_0) &\left\{\prod_{k=1}^{\mu-1} d(\alpha_{k+1}-\alpha_k)\right\} d(\alpha_1-\alpha'_0) \left\{\prod_{k=1}^{\mu-1} I(\alpha_{k+1}-\alpha'_k)\right\} I(\alpha_1-\alpha_0), \end{aligned} \tag{5.34}$$

where we have used the relation :

$$\frac{c(\alpha'_1-\alpha_1)}{b(\alpha'_1-\alpha_1)}e^{\rho(\alpha'_1-\alpha_1)}-\frac{c(\alpha_0-\alpha'_0)}{b(\alpha_0-\alpha'_0)}e^{\rho(\alpha_0-\alpha'_0)} = -\frac{\mathrm{sh}\left(\rho(\frac{2\pi i}{n})\right)\mathrm{sh}\left(\rho(\alpha_0+\alpha_1-\alpha'_0-\alpha'_1))\right)}{\mathrm{sh}\left(\rho(\alpha'_1-\alpha_1)\right)\mathrm{sh}\left(\rho(\alpha_0-\alpha'_0))\right)}. \tag{5.35}$$

We arrive at

$$A_\mu+B_\mu+C_\mu \sim A_\mu+A''_\mu+C_\mu+C'_\mu \sim 0. \tag{5.36}$$

Now we have proved the commutation relation (3.1). As the same arguments as the above we can show the commutation relations (3.2), (3.4) and (3.10).

Next we prove the inversion relations.

The bosonization of the type I vertex operator is deformed as

$$\begin{aligned}\Phi_j(\beta) &= \left(\frac{i}{\pi}\right)^j e^{-j\frac{\xi-1}{\xi}(\gamma+\log\frac{2\pi\xi}{n})}\int_{C_j}\frac{d\alpha_j}{2\pi i}\cdots\int_{C_1}\frac{d\alpha_1}{2\pi i}e^{\frac{n}{2\xi}(\alpha_j-\beta)} \\ &\times\ :U_0(\beta)\cdots U_j(\alpha_j):\prod_{k=1}^{j}\Gamma\left(\frac{n}{2\pi\xi}i(\alpha_k-\alpha_{k-1})+\frac{1}{2\xi}\right)\Gamma\left(-\frac{n}{2\pi\xi}i(\alpha_k-\alpha_{k-1})+\frac{1}{2\xi}\right),\end{aligned} \tag{5.37}$$

where $\alpha_0=\beta$. Here the contour $C_k,\ (k=1,\cdots,j)$ is taken $(-\infty,\infty)$ except that the poles

$$\alpha_k-\alpha_{k-1}=\frac{\pi i}{n}+\frac{2\pi\xi i}{n}l,\quad (l\in\mathbb{N}\geq 0), \tag{5.38}$$

of $\Gamma\left(\frac{n}{2\pi\xi}i(\alpha_k-\alpha_{k-1})+\frac{1}{2\xi}\right)$ are above C_k and the poles

$$\alpha_k-\alpha_{k-1}=-\frac{\pi i}{n}-\frac{2\pi\xi i}{n}l,\quad (l\in\mathbb{N}\geq 0), \tag{5.39}$$

of $\Gamma\left(-\frac{n}{2\pi\xi}i(\alpha_k-\alpha_{k-1})+\frac{1}{2\xi}\right)$ are below C_k.

The bosonization of the dual type-I vertex operator is deformed as

$$\begin{aligned}\Phi_j^*(\beta) &= \left(\frac{i}{\pi}\right)^{n-j-1} e^{-(n-j-1)\frac{\xi-1}{\xi}(\gamma+\log\frac{2\pi\xi}{n})}\int_{C_{j+1}^*}\frac{d\alpha_{j+1}}{2\pi i}\cdots\int_{C_{n-1}^*}\frac{d\alpha_{n-1}}{2\pi i}e^{\frac{n}{2\xi}(\alpha_{j+1}-\beta)} \\ &\times\ :U_{j+1}(\alpha_{j+1})\cdots U_n(\beta):\prod_{k=j+1}^{n-1}\Gamma\left(\frac{n}{2\pi\xi}i(\alpha_{k+1}-\alpha_k)+\frac{1}{2\xi}\right)\Gamma\left(-\frac{n}{2\pi\xi}i(\alpha_{k+1}-\alpha_k)+\frac{1}{2\xi}\right),\end{aligned} \tag{5.40}$$

where $\alpha_n=\beta$. Here the contour $C_k^*,\ (k=j+1,\cdots,n-1)$ is taken $(-\infty,\infty)$ except that the poles

$$\alpha_k-\alpha_{k+1}=\frac{\pi i}{n}+\frac{2\pi\xi i}{n}l,\quad (l\in\mathbb{N}\geq 0), \tag{5.41}$$

of $\Gamma\left(-\frac{n}{2\pi\xi}i(\alpha_{k+1}-\alpha_k)+\frac{1}{2\xi}\right)$ are above C_k^* and the poles

$$\alpha_k-\alpha_{k+1}=-\frac{\pi i}{n}-\frac{2\pi\xi i}{n}l,\quad (l\in\mathbb{N}\geq 0), \tag{5.42}$$

of $\Gamma\left(\frac{n}{2\pi\xi}i(\alpha_{k+1}-\alpha_k)+\frac{1}{2\xi}\right)$ are below C_k^*.

Let us prove the relation (3.12). For $j_1<j_2$ we have

$$\Phi_{j_1}(\beta_1)\Phi_{j_2}^*(\beta_2)=r^*(\beta_1-\beta_2)\Phi_{j_2}^*(\beta_2)\Phi_{j_1}(\beta_1). \tag{5.43}$$

As $\beta_2\to\beta_1+\pi i$, $r^*(\beta_1-\beta_2)$ becomes zero.

Next we prove the case $j_1=j_2$. We have

$$\begin{aligned}\Phi_j(\beta_1)\Phi_j^*(\beta_2) &= r^*(\beta_1-\beta_2)\int_{C_1}\frac{d\alpha_1}{2\pi i}\cdots\int_{C_j}\frac{d\alpha_j}{2\pi i}\int_{C_{j+1}^*}\frac{d\alpha_{j+1}}{2\pi i}\cdots\int_{C_{n-1}^*}\frac{d\alpha_{n-1}}{2\pi i} \\ &\times\ U_n(\beta_2)U_{n-1}(\alpha_{n-1})\cdots U_1(\alpha_1)U_0(\beta_1) \\ &\times\ e^{\frac{n}{2\xi}(\alpha_j+\alpha_{j+1}-\beta_1-\beta_2)}\mathrm{sh}\left(\frac{n}{2\xi}(\alpha_j-\alpha_{j+1}-\frac{\pi i}{n})\right)\prod_{k=1}^{n}\frac{1}{\mathrm{sh}\left(\frac{n}{2\xi}(\alpha_k-\alpha_{k-1}-\frac{\pi i}{n})\right)}.\end{aligned} \tag{5.44}$$

When we take the limit $\beta_2 \to \beta_1 + \pi i$, the contour is pinched but the function $r^*(\beta_1 - \beta_2)$ has a zero. Therefore the limit is evaluated by successively taking the residues at $\alpha_k = \alpha_{k-1} + \frac{\pi i}{n}$ for $k = 1, \cdots, j$, and $\alpha_k = \alpha_{k+1} - \frac{\pi i}{n}$ for $k = j+1, \cdots, n-1$, successively. The following relation is usefull.

$$U_n(\beta + \pi i) U_{n-1}\left(\beta + \frac{n-1}{n}\pi i\right) \cdots U_1\left(\beta + \frac{\pi i}{n}\right) U_0(\beta) = e^{-\frac{\xi-1}{\xi}(\gamma + \log\frac{2\pi\xi}{n})n}\Gamma(1/\xi)^n. \tag{5.45}$$

Now we have proved the inversion relation (3.12).

6 Correlation functions

In this section we derive a solution of the system of difference equations (1.3) and (1.4), algebraically, and obtain an integral representation of it.

Let us introduce the degree operator D on the Fock space :

$$D\, b_j(-t)|0\rangle = t\, b_j(-t)|0\rangle, \ (t > 0). \tag{6.1}$$

We have

$$e^{\lambda D} b(t) e^{-\lambda D} = e^{-\lambda t} b(t). \tag{6.2}$$

Therefore the degree operator D has the homogeneity condition.

$$e^{\lambda D} U_j(\beta) e^{-\lambda D} = U_j(\beta + i\lambda). \tag{6.3}$$

The vertex operator and the degree operator enjoy the homogeneity property.

$$e^{-\lambda D}\Phi_j(\beta) e^{\lambda D} = \ \Phi_j(\beta + i\lambda), \quad e^{-\lambda D}\Phi_j^*(\beta) e^{\lambda D} = \Phi_j^*(\beta + i\lambda). \tag{6.4}$$

Now let us consider the trace functions for $\lambda > 0$ defined by

$$\begin{aligned} & G(\beta_1 \cdots \beta_N | \beta_{N+1} \cdots \beta_{2N})_{\epsilon_1 \cdots \epsilon_{2N}} \\ = \ & \frac{\mathrm{tr}_{\mathcal{H}}\left(e^{-\lambda D}\Phi^*_{\epsilon_1}(\beta_1) \cdots \Phi^*_{\epsilon_N}(\beta_N)\Phi_{\epsilon_{N+1}}(\beta_{N+1}) \cdots \Phi_{\epsilon_{2N}}(\beta_{2N})\right)}{\mathrm{tr}_{\mathcal{H}}\left(e^{-\lambda D}\right)}, \end{aligned} \tag{6.5}$$

where the space $\mathcal{H}$ is the Fock space of the free bosons.

By using the homogeneity condition (6.4) and the commutation relations (3.1), (3.2), (3.10), it is shown that the above trace function satisfies the desired difference equations (1.3), (1.4). The inversion relation (3.12) means the restriction equation (1.5).

Using the free boson realizations, we shall treat the trace of the form : $\mathrm{tr}_{\mathcal{H}}(e^{-\lambda D}\mathcal{O})$. The calculation is simplified by the technique of Clavelli and Shapiro [6]. Their prescription is as follows. Introduce a copy of the bosons $a(t)$ $(t \in \mathbb{R})$ satisfying the relation $[a(t), b(t')] = 0$ and the same commutation relation as the $b(t)$. Let

$$\tilde{b}(t) = \frac{b(t)}{1 - e^{-\lambda t}} + a(-t), \quad \tilde{b}(-t) = \frac{a(t)}{e^{\lambda t} - 1} + b(-t), \quad (t > 0). \tag{6.6}$$

For a linear operator $\mathcal{O}$ on the Fock space $\mathcal{H}_b = \mathcal{F}[b(-t)]$, let $\tilde{\mathcal{O}}$ be the operator on $\mathcal{H}_b \otimes \mathcal{H}_a$, ($\mathcal{H}_a = \mathcal{F}[a(-t)]$) obtaind by substituting $\tilde{b}(t)$ for $b(t)$. We have then

$$\frac{\mathrm{tr}_{\mathcal{H}_b}\left(e^{-\lambda D}\mathcal{O}\right)}{\mathrm{tr}_{\mathcal{H}_b}(e^{-\lambda D})} = \langle\tilde{0}|\tilde{\mathcal{O}}|\tilde{0}\rangle, \tag{6.7}$$

where the left hand side denotes the usual expectation value with respect to the Fock vacuum $|\tilde{0}\rangle = |0\rangle \otimes |0\rangle$, $\langle\tilde{0}|\tilde{0}\rangle$. In what follows we use the following abbreviation :

$$\langle\mathcal{O}\rangle_\lambda = \frac{\mathrm{tr}_{\mathcal{H}}\left(e^{-\lambda D}\mathcal{O}\right)}{\mathrm{tr}_{\mathcal{H}}(e^{-\lambda D})}. \tag{6.8}$$

We have the following.

$$\langle b_j(t)b_k(t')\rangle_\lambda = \frac{e^{\lambda t}}{e^{\lambda t}-1}[b_j(t), b_k(t')]. \tag{6.9}$$

We have the following formula usefull to evaluate the function (6.5).

$$\frac{\mathrm{tr}_H\left(e^{-\lambda D} : e^{\int_{-\infty}^{\infty} c(t)e^{i\beta_1 t}dt} :: e^{\int_{-\infty}^{\infty} d(t)e^{i\beta_1 t}dt} :\right)}{\mathrm{tr}_H(e^{-\lambda D})} = \exp\left(\int_0^\infty A(t)\frac{\mathrm{ch}(i(\beta_1-\beta_2)+\frac{\lambda}{2})t}{\mathrm{sh}\frac{\lambda t}{2}}dt\right), \tag{6.10}$$

where $c(t)$ and $d(t)$ are bosons satisfying $[c(t), d(t')] = A(t)\delta(t+t')$ and $A(t) = -A(-t)$. In order to understand an integral of the right hand side, see the Appendix B.

The basic trace functions are evaluated as following.

$$\langle U_0(\alpha_1)U_0(\alpha_2)\rangle_\lambda = \langle U_n(\alpha_1)U_n(\alpha_2)\rangle_\lambda = \mathrm{Const}.E_\lambda(\alpha_1-\alpha_2), \tag{6.11}$$

$$\langle U_0(\alpha_1)U_n(\alpha_2)\rangle_\lambda = \langle U_n(\alpha_1)U_0(\alpha_2)\rangle_\lambda = \mathrm{Const}.E^*_\lambda(\alpha_1-\alpha_2). \tag{6.12}$$

Here we set

$$E_\lambda(\alpha) = \frac{S_3(-i\alpha+\frac{2\pi}{n})S_3(i\alpha+\frac{2\pi}{n}+\lambda)}{S_3(-i\alpha+\frac{2\pi}{n}\xi)S_3(i\alpha+\frac{2\pi}{n}\xi+\lambda)} \tag{6.13}$$

$$E^*_\lambda(\alpha) = \frac{S_3(-i\alpha+\pi)S_3(i\alpha+\lambda+\pi)}{S_3(-i\alpha+\frac{2\pi}{n}+\pi)S_3(i\alpha+\lambda+\pi+\frac{2\pi}{n})}, \tag{6.14}$$

where

$$S_3(\beta) = S_3\left(\beta\middle|2\pi, \frac{2\pi}{n}\xi, \lambda\right). \tag{6.15}$$

$$\begin{aligned}&\langle U_j(\alpha_1)U_{j-1}(\alpha_2)\rangle_\lambda = \langle U_{j-1}(\alpha_1)U_j(\alpha_2)\rangle_\lambda\\ = \quad &\mathrm{Const}.\varphi(\alpha_1-\alpha_2)\times\frac{1}{\mathrm{sh}\left(\frac{n}{2\xi}(\alpha_1-\alpha_2-\frac{\pi i}{n}+\lambda i)\right)},\end{aligned} \tag{6.16}$$

$$\begin{aligned}&\langle U_j(\alpha_1)U_j(\alpha_2)\rangle_\lambda = \mathrm{Const}.\psi(\alpha_1-\alpha_2)\times\\ \times\quad &\mathrm{sh}\left(\frac{n}{2\xi}\alpha\right)\mathrm{sh}\left(\frac{n}{2\xi}(\alpha+\frac{2\pi i}{n})\right)\mathrm{sh}\left(\frac{\pi}{\lambda}(\alpha+\frac{2\pi i}{n})\right)\mathrm{sh}\left(\frac{\pi}{\lambda}(-\alpha+\frac{2\pi i}{n})\right).\end{aligned} \tag{6.17}$$

Here we set

$$\varphi(\alpha) = \frac{1}{S_2\left(i\alpha+\frac{\pi}{n}\Big|\lambda,\frac{2\pi}{n}\xi\right)S_2\left(-i\alpha+\frac{\pi}{n}\Big|\lambda,\frac{2\pi}{n}\xi\right)}, \tag{6.18}$$

$$\psi(\alpha) = \frac{1}{S_2\left(i\alpha-\frac{2\pi}{n}\Big|\lambda,\frac{2\pi}{n}\xi\right)S_2\left(-i\alpha-\frac{2\pi}{n}\Big|\lambda,\frac{2\pi}{n}\xi\right)}. \tag{6.19}$$

The functions $\varphi(\alpha)$ and $\psi(\alpha)$ become the integral kernel of the correlation functions.
The function $\varphi(\alpha)$ has poles at

$$\alpha = \pm i\left(n_1\lambda+n_2\frac{2\pi}{n}\xi+\frac{\pi}{n}\right), \quad (n_1,n_2\geq 0). \tag{6.20}$$

The function $\psi(\alpha)$ has poles at

$$\alpha = \pm i\left(n_1\lambda+n_2\frac{2\pi}{n}\xi-\frac{2\pi}{n}\right), \quad (n_1,n_2\geq 0). \tag{6.21}$$

The trace of the vertex operators (6.5) is evaluated by applying the Wick's theorem.
The one-point correlation functions are evaluated as follows. In what follows we set $\rho=\frac{n}{2\xi}$.

$$\begin{aligned} G(\beta_1|\beta_1')_{\epsilon,\epsilon} &= E_\lambda^*(\beta_1-\beta_1')\int_{-\infty}^{\infty}\frac{d\alpha_1}{2\pi i}\cdots\int_{-\infty}^{\infty}\frac{d\alpha_{n-1}}{2\pi i}\prod_{k=1}^{n}\varphi(\alpha_k-\alpha_{k-1})e^{\rho(\alpha_\epsilon+\alpha_{\epsilon+1}-\beta_1-\beta_1')} \\ &\times \ \mathrm{sh}\left(\rho(\alpha_{\epsilon+1}-\alpha_\epsilon-\frac{\pi i}{n})\right)\prod_{k=1}^{n}\frac{1}{\mathrm{sh}\left(\rho(\alpha_k-\alpha_{k-1}-\frac{\pi i}{n})\right)\mathrm{sh}\left(\rho(\alpha_k-\alpha_{k-1}-\frac{\pi i}{n}+\lambda i)\right)}, \end{aligned} \tag{6.22}$$

where $\alpha_n=\beta_1,\ \alpha_0=\beta_1'$. Here we omit an irrelevant constant factor.
The N point correlation functions are evaluated as follows.

$$G(\beta_1\cdots\beta_N|\beta_N'\cdots\beta_1')_{\epsilon_1\cdots\epsilon_N,\epsilon_N,\cdots,\epsilon_1} = E(\beta_1\cdots\beta_N|\beta_N'\cdots\beta_1')\prod_{j,r}\int_{-\infty}^{\infty}\frac{d\alpha_{j,r}}{2\pi i}I_{\epsilon_1\cdots\epsilon_N}(\{\alpha_{j,r}\}). \tag{6.23}$$

We associate the variables $\alpha_{j,r},\ (1\leq r\leq N, 0\leq j\leq \epsilon_r)$ to the basic operator $U_j(\alpha_{j,r})$ contained in the vertex operators $\Phi_{\epsilon_r}(\beta_r')$ and the variables $\alpha_{j,r},\ (1\leq r\leq N, \epsilon_r+1\leq j\leq n)$ to the basic operator $U_j(\alpha_{j,r})$ contained in the vertex operators $\Phi_{\epsilon_r}^*(\beta_r)$.
We set

$$\alpha_{n,r}=\beta_r, \qquad \alpha_{0,r}=\beta_r', \tag{6.24}$$

$$\mathcal{N}_j^*=\{k|\epsilon_k\leq j-1\}, \qquad \mathcal{N}_j=\{k|\epsilon_k\geq j\}. \tag{6.25}$$

The function $E(\beta_1\cdots\beta_N|\beta_N'\cdots\beta_1')$ is given by

$$\begin{aligned} &E(\beta_1\cdots\beta_N|\beta_N'\cdots\beta_1') \\ =\ & e^{-\rho(\beta_1+\cdots+\beta_N+\beta_1'+\cdots+\beta_N')}\prod_{1\leq j<k\leq N}E_\lambda(\beta_j-\beta_k)\prod_{1\leq j<k\leq N}E_\lambda(\beta_k'-\beta_j')\prod_{j,k=1}^{N}E_\lambda^*(\beta_j-\beta_k'). \end{aligned} \tag{6.26}$$

The integrand is given by

$$I_{\epsilon_1\cdots\epsilon_N}(\{\alpha_{j,r}\}) = K(\{\alpha_{j,r}\})g(\{\alpha_{j,r}\})h(\{\alpha_{j,r}\}). \tag{6.27}$$

Here the integral kernel is given by

$$K(\{\alpha_{j,r}\}) = \prod_{j=1}^{n-1} \prod_{\substack{r,s=1 \\ r>s}}^{N} \psi(\alpha_{j,r}-\alpha_{j,s}) \times \prod_{j=1}^{n-1} \prod_{k,l=1}^{N} \left\{ \prod_{\substack{k\in N_j^* \\ l\in N_{j-1}}} \varphi(\alpha_{j,k}-\alpha_{j,l}) \prod_{\substack{k\in N_{j-1}^* \\ l\in N_j}} \varphi(\alpha_{j,k}-\alpha_{j,l}) \right\}$$
$$\times \prod_{j=1}^{n-1} \prod_{k,l=1}^{N} \left\{ \prod_{\substack{k\in N_j \\ l\in N_{j-1}}} \varphi(\alpha_{j,k}-\alpha_{j,l}) \prod_{\substack{k\in N_j^* \\ l\in N_{j-1}^*}} \varphi(\alpha_{j,k}-\alpha_{j,l}) \right\}^2 . \quad (6.28)$$

Here we set

$$h(\{\alpha_{j,r}\}) = \prod_{j=1}^{n-1} \prod_{\substack{r,s=1 \\ r>s}}^{N} \left\{ \mathrm{sh}\left(\frac{\pi}{\lambda}(\alpha_{j,r}-\alpha_{j,s}+\frac{2\pi i}{n})\right) \mathrm{sh}\left(\frac{\pi}{\lambda}(\alpha_{j,s}-\alpha_{j,r}+\frac{2\pi i}{n})\right) \right\}, \quad (6.29)$$

and

$$\begin{aligned}
& g(\{\alpha_{j,r}\}) \\
= \; & \prod_{j=1}^{n-1} \prod_{\substack{r>s \\ s\in N_j}} \left\{ \mathrm{sh}\left(\rho(\alpha_{j,r}-\alpha_{j,s})\right) \mathrm{sh}\left(\rho(\alpha_{j,r}-\alpha_{j,s}+\frac{2\pi i}{n})\right) \right\} \\
\times \; & \prod_{j=1}^{n-1} \prod_{\substack{r>s \\ s\in N_j^*}} \left\{ \mathrm{sh}\left(\rho(\alpha_{j,s}-\alpha_{j,r})\right) \mathrm{sh}\left(\rho(\alpha_{j,s}-\alpha_{j,r}+\frac{2\pi i}{n})\right) \right\} \\
\times \; & \prod_{j=1}^{n} \prod_{\substack{r\in N_j \\ s\in N_{j-1}}} \left\{ \mathrm{sh}\left(\rho(\alpha_{j,r}-\alpha_{j-1,s}-\frac{\pi i}{n}+\lambda i)\right) \mathrm{sh}\left(\rho(\alpha_{j-1,r}-\alpha_{j,s}-\frac{\pi i}{n}+\lambda i)\right) \right\}^{-1} \\
\times \; & \prod_{j=1}^{n} \prod_{\substack{r\in N_j^* \\ s\in N_{j-1}^*}} \left\{ \mathrm{sh}\left(\rho(\alpha_{j,r}-\alpha_{j-1,s}-\frac{\pi i}{n}+\lambda i)\right) \mathrm{sh}\left(\rho(\alpha_{j-1,r}-\alpha_{j,s}-\frac{\pi i}{n}+\lambda i)\right) \right\}^{-1} \qquad (6.30) \\
\times \; & \prod_{j=1}^{n} \left\{ \prod_{\substack{r\in N_j^* \\ s\in N_{j-1}}} \mathrm{sh}\left(\rho(\alpha_{j,r}-\alpha_{j-1,s}-\frac{\pi i}{n}+\lambda i)\right) \prod_{\substack{r\in N_{j-1}^* \\ s\in N_j}} \mathrm{sh}\left(\rho(\alpha_{j,r}-\alpha_{j-1,s}-\frac{\pi i}{n}+\lambda i)\right) \right\}^{-1} \\
\times \; & \prod_{r=1}^{N} e^{\rho(\alpha_{\epsilon_r}+\alpha_{\epsilon_r+1})} \prod_{r=1}^{N} \mathrm{sh}\left(\rho(\alpha_{\epsilon_r+1,r}-\alpha_{\epsilon_r,r}-\frac{\pi i}{n})\right) \prod_{j=1}^{n} \prod_{r=1}^{N} \left\{ \mathrm{sh}\left(\rho(\alpha_{j,r}-\alpha_{j-1,r}-\frac{\pi i}{n})\right) \right\}^{-1} .
\end{aligned}$$

Here we omit an irrelevant constant factor.

Acknowledgements. This work was partly supported by Grant-in-Aid for Encouragements for Young Scientists (A) from Japan Society for the Promotion of Science. (11740099)

References

[1] M.Jimbo,H.Konno and T.Miwa: Massless XXZ model and degeneration of the elliptic algebra $\mathcal{A}_{q,p}(\widehat{sl_2})$, *Deformation theory and Symplectic Geometry*, Kluwere Academic Publishers, (1997).

[2] M.Jimbo and T.Miwa: Quantum KZ equation with $|q|=1$ and correlation functions of the XXZ model in the gapless regime, *J.Phys.***A29**,2923-2958,(1996).

[3] T. Miwa and Y. Takeyama : The integral formula for the solutions of the quantum Knizhnik-Zamolodchikov equation associated with $U_q(\widehat{sl_n})$ for $|q|=1$, *Proceedings for SIDE III meeting at Sabaudia near Rome*, (1999).

[4] Y. Koyama : Staggered Polarization of Vertex Models with $U_q(\widehat{sl_n})-$Symmetry, *Commun. Math. Phys.***164**, 277-291, (1994).

[5] A.Doikou and R.I.Nepomechie: Soliton S matrices for the critical $A_{N-1}^{(1)}$ chain, *Phys. Lett.***B462**,127-131, (1999).

[6] L.Clavelli and J.A.Shapiro : Pomeron Factorization in General Dual Models, *Nucl.Phys.***B57**, 490-535, (1973).

[7] Y.Asai, M.Jimbo, T.Miwa and Ya. Pugai : Bosonization of Vertex Operators for $A_{n-1}^{(1)}$ Face Model, *J.Phys.***A29**, 6596-6616, (1996).

[8] H. Furutsu, T.Kojima and Y.-H. Quano : Type II Vertex Operators for the $A_{n-1}^{(1)}$ Face Model, to appear in *Int.J.Mod.Phys.***A**.

[9] N. Kurokawa : On the generalization of the sine function, *Technical Report of Tsuda University,* 4:1-25, (1992).

[10] S. Lukyanov : Free field representation for massive integrable models, *Commun.Math.Phys.***167**, 183-226, (1995).

[11] F. Smirnov : Form Factors in Completely Integrable Models of Quantum Field Theory, Advanced Series in Mathematical Physics Vol.14, World Scientific, 1992.

A Multi-Gamma functions

Here we summarize the multiple gamma and the multiple sine functions, following Kurokawa [9].
Let us set the functions $\Gamma_1(x|\omega), \Gamma_2(x|\omega_1,\omega_2)$ and $\Gamma_3(x|\omega_1,\omega_2,\omega_3)$ by

$$\log\Gamma_1(x|\omega)+\gamma B_{11}(x|\omega) = \int_C \frac{dt}{2\pi i t}e^{-xt}\frac{\log(-t)}{1-e^{-\omega t}}, \tag{A.1}$$

$$\log\Gamma_2(x|\omega_1,\omega_2)-\frac{\gamma}{2}B_{22}(x|\omega_1,\omega_2) = \int_C \frac{dt}{2\pi i t}e^{-xt}\frac{\log(-t)}{(1-e^{-\omega_1 t})(1-e^{-\omega_2 t})}, \tag{A.2}$$

$$\log\Gamma_3(x|\omega_1,\omega_2,\omega_3)+\frac{\gamma}{3!}B_{33}(x|\omega_1,\omega_2,\omega_3) = \int_C \frac{dt}{2\pi i t}e^{-xt}\frac{\log(-t)}{(1-e^{-\omega_1 t})(1-e^{-\omega_2 t})(1-e^{-\omega_3 t})}, \tag{A.3}$$

where the functions $B_{jj}(x)$ are the multiple Bernoulli polynomials defined by

$$\frac{t^r e^{xt}}{\prod_{j=1}^r (e^{\omega_j t}-1)} = \sum_{n=0}^\infty \frac{t^n}{n!}B_{r,n}(x|\omega_1\cdots\omega_r), \tag{A.4}$$

more explicitly

$$B_{11}(x|\omega) = \frac{x}{\omega} - \frac{1}{2}, \tag{A.5}$$

$$B_{22}(x|\omega) = \frac{x^2}{\omega_1\omega_2} - \left(\frac{1}{\omega_1} + \frac{1}{\omega_2}\right)x + \frac{1}{2} + \frac{1}{6}\left(\frac{\omega_1}{\omega_2} + \frac{\omega_2}{\omega_1}\right). \tag{A.6}$$

Here γ is Euler's constant, $\gamma = \lim_{n\to\infty}(1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \log n)$.
Here the contor of integral is given by



Contour C

Let us set

$$S_1(x|\omega) = \frac{1}{\Gamma_1(\omega - x|\omega)\Gamma_1(x|\omega)}, \tag{A.7}$$

$$S_2(x|\omega_1,\omega_2) = \frac{\Gamma_2(\omega_1 + \omega_2 - x|\omega_1,\omega_2)}{\Gamma_2(x|\omega_1,\omega_2)}, \tag{A.8}$$

$$S_3(x|\omega_1,\omega_2,\omega_3) = \frac{1}{\Gamma_3(\omega_1 + \omega_2 + \omega_3 - x|\omega_1,\omega_2,\omega_3)\Gamma_3(x|\omega_1,\omega_2,\omega_3)} \tag{A.9}$$

We have

$$\Gamma_1(x|\omega) = e^{(\frac{x}{\omega} - \frac{1}{2})\log\omega}\frac{\Gamma(x/\omega)}{\sqrt{2\pi}}, \ S_1(x|\omega) = 2\sin(\pi x/\omega), \tag{A.10}$$

$$\frac{\Gamma_2(x + \omega_1|\omega_1,\omega_2)}{\Gamma_2(x|\omega_1,\omega_2)} = \frac{1}{\Gamma_1(x|\omega_2)}, \ \frac{S_2(x + \omega_1|\omega_1,\omega_2)}{S_2(x|\omega_1,\omega_2)} = \frac{1}{S_1(x|\omega_2)}, \ \frac{\Gamma_1(x + \omega|\omega)}{\Gamma_1(x|\omega)} = x. \tag{A.11}$$

$$\frac{\Gamma_3(x + \omega_1|\omega_1,\omega_2,\omega_3}{\Gamma_3(x|\omega_1,\omega_2,\omega_3)} = \frac{1}{\Gamma_2(x|\omega_2,\omega_3)}, \ \frac{S_3(x + \omega_1|\omega_1,\omega_2,\omega_3)}{S_3(x|\omega_1,\omega_2,\omega_3)} = \frac{1}{S_2(x|\omega_2,\omega_3)}. \tag{A.12}$$

$$\log S_2(x|\omega_1\omega_2) = \int_C \frac{\mathrm{sh}(x - \frac{\omega_1+\omega_2}{2})t}{2\mathrm{sh}\frac{\omega_1 t}{2}\mathrm{sh}\frac{\omega_2 t}{2}}\log(-t)\frac{dt}{2\pi i t}, \ (0 < \mathrm{Re}x < \omega_1 + \omega_2). \tag{A.13}$$

$$S_2(x|\omega_1\omega_2) = \frac{2\pi}{\sqrt{\omega_1\omega_2}}x + O(x^2), \quad (x \to 0). \tag{A.14}$$

$$S_2(x|\omega_1\omega_2)S_2(-x|\omega_1\omega_2) = -4\sin\frac{\pi x}{\omega_1}\sin\frac{\pi x}{\omega_2}. \tag{A.15}$$

B Normal Ordering

Here we list the formulas of the form

$$X(\beta_1)Y(\beta_2) = C_{XY}(\beta_1 - \beta_2) : X(\beta_1)X(\beta_2) :, \tag{B.1}$$

where $X, Y = U_j$, and $C_{XY}(\beta)$ is a meromorphic function on $\mathbb{C}$. These formulae follow from the commutation relation of the free bosons. When we compute the contraction of the basic operators, we often encounter an integral

$$\int_0^\infty F(t)dt, \tag{B.2}$$

which is divergent at $t = 0$. Here we adopt the following prescription for regularization : it should be understood as the countour integral,

$$\int_C F(t)\frac{\log(-t)}{2\pi i}dt, \tag{B.3}$$

where the countour C is given by



Contour C

The contractions of the basic operators have the following forms.

$$U_j(\alpha_1)U_j(\alpha_2) = h_{jj}(\alpha_1 - \alpha_2) : U_j(\alpha_1)U_j(\alpha_2) : \quad \left(\mathrm{Im}(\alpha_2 - \alpha_1) < \frac{2\pi}{n},\ j = 0, n\right), \tag{B.4}$$

$$h_{00}(\alpha) = h_{nn}(\alpha) = e^{\gamma\frac{\xi-1}{\xi}\frac{n-1}{n}} \frac{\Gamma_2(-i\alpha + \frac{2\pi}{n}\xi|\frac{2\pi}{n}\xi, 2\pi)\Gamma_2(-i\alpha + 2\pi|\frac{2\pi}{n}\xi, 2\pi)}{\Gamma_2(-i\alpha + \frac{2\pi}{n}|\frac{2\pi}{n}\xi, 2\pi)\Gamma_2(-i\alpha + 2\pi + \frac{2\pi}{n}\xi - \frac{2\pi}{n}|\frac{2\pi}{n}\xi, 2\pi)}, \tag{B.5}$$

$$U_j(\alpha_1)U_j(\alpha_2) = h_{jj}(\alpha_1 - \alpha_2) : U_j(\alpha_1)U_j(\alpha_2) : \quad \left(\mathrm{Im}(\alpha_2 - \alpha_1) < \frac{2\pi}{n}(\xi - 1),\ 1 \le j \le n - 1\right) \tag{B.6}$$

$$h_{jj}(\alpha) = e^{2\gamma\frac{\xi-1}{\xi}} \frac{\alpha}{i} \frac{\Gamma_1(-i\alpha + \frac{2\pi}{n}\xi - \frac{2\pi}{n}|\frac{2\pi}{n}\xi)}{\Gamma_1(-i\alpha + \frac{2\pi}{n}|\frac{2\pi}{n}\xi)}, \tag{B.7}$$

$$U_j(\alpha_1)U_{j-1}(\alpha_2) = h_{jj-1}(\alpha_1 - \alpha_2) : U_{j-1}(\alpha_1)U_j(\alpha_2) : \quad \left(\mathrm{Im}(\alpha_2 - \alpha_1) < \frac{\pi}{n},\ 1 \le j \le n\right), \tag{B.8}$$

$$h_{jj-1}(\alpha) = e^{-\gamma\frac{\xi-1}{\xi}} \frac{\Gamma_1(-i\alpha + \frac{\pi}{n}|\frac{2\pi}{n}\xi)}{\Gamma_1(-i\alpha + \frac{\pi}{n}(2\xi - 1)|\frac{2\pi}{n}\xi)}, \tag{B.9}$$

and

$$U_0(\alpha_1)U_n(\alpha_2) = h_{0n}(\alpha_1 - \alpha_2) : U_0(\alpha_1)U_n(\alpha_2) :, \quad \left(\mathrm{Im}(\alpha_2 - \alpha_1) < \frac{2\pi}{n}\right) \tag{B.10}$$

$$h_{0n}(\alpha) = e^{\frac{\gamma}{n}\frac{\xi-1}{\xi}} \frac{\Gamma_2(-i\alpha + \pi + \frac{2\pi}{n}\xi - \frac{2\pi}{n}|\frac{2\pi}{n}\xi, 2\pi)\Gamma_2(-i\alpha + \pi + \frac{2\pi}{n}|\frac{2\pi}{n}\xi, 2\pi)}{\Gamma_2(-i\alpha + \pi|\frac{2\pi}{n}\xi, 2\pi)\Gamma_2(-i\alpha + \pi + \frac{2\pi}{n}\xi|\frac{2\pi}{n}\xi, 2\pi)}, \tag{B.11}$$